

1. <u>MMR</u> (Measles. Mumps. and Rubella): If born after January 1957, 2 doses of MMR 28 days apart or serological (blood test) evidence of immunity.

2. <u>Chicken Pox</u> (Varicella): 2 vaccines given at least 4 weeks apart are required. Documentation of the disease or serological proof of immunity is also accepted.

3. <u>Tdap</u> (Tetanus. Diphtheria. Pertussis): A booster of "Tdap" is required within the past 10 years.

4. <u>Hepatitis B Series</u>: 3 vaccinations or serological evidence of immunity. The administration dosage and time separation of the 3 Hepatitis B vaccination series as recommended by the Texas Health and Human Services. For previously unvaccinated or incompletely vaccinated adults 18 years of age or older, Heplisav-B (HepB-CpG) 2 dose vaccination administered 4 weeks apart will be accepted.

5. Flu Vaccine: Within the past 12 months unless the vaccine is not available.

6. <u>Meningococcal</u> (MCV4): If under the age of 22 (Vaccine received within the last 5 years of enrollment).

7. <u>**TB Skin Test**</u>: within the past 12 months. If the skin test is positive, a negative chest x-ray or official documentation of negative chest x-ray is required.

8. <u>COVID-19</u>: One-dose Johnson & Johnson Janssen Vaccine (May receive booster at least 2 months after primary dose)

-----or either brand below-----

Two-dose Moderna Vaccine given 28 days apart (May receive booster 6 months after last dose) Two-dose Pfizer Vaccine given 21 days apart (May receive booster 6 months after last dose)